arXiv:1603.05728v1 [math.CV] 17 Mar 2016

LELONG NUMBERS, COMPLEX SINGULARITY EXPONENTS, AND SIU'S SEMICONTINUITY THEOREM

QI'AN GUAN AND XIANGYU ZHOU

Abstract. In this note, we present a relationship between Lelong numbers and complex singularity exponents. As an application, we obtain a new proof of Siu's semicontinuity theorem for Lelong numbers.

RÉSUMÉ. Dans cette note, nous présentons une relation entre le nombre de Lelong et exposants de singularités complexes. Comme application, nous obtenons une nouvelle preuve de semicontinuité théorème de Siu pour les numéros Lelong.

1. Introduction

Let φ be a plurisubharmonic function near the origin $o \in \mathbb{C}^n$. The Lelong number is defined as follows

Definition 1.1. $\nu(\varphi, o) := \sup\{c \geq 0 : \varphi \leq c \log|z| + O(1)\}$.

In [19], Siu established the semicontinuity theorem for Lelong numbers, i.e., the upper level set of Lelong numbers is analytic. After that, Kiselman [15] generalized Siu's semicontinuity theorem for directed Lelong numbers. In [2], Demailly introduced a generalized Lelong number and generalized the above result of Kiselman. And then, Demailly (see [5]) gave a completely new and simple proof of Siu's theorem by using Ohsawa-Takegoshi L^2 extension theorem.

In this note, we present a new proof of Siu's theorem by establishing a relationship between Lelong numbers and complex singularity exponents.

Now let us recall the definition of the complex singularity exponent using the concept of multiplier ideal sheaf $\mathcal{I}(\varphi)$ (see [22], see also [3, 5]) (also called log canonical threshold in algebraic geometry see [18, 17]):

Definition 1.2. *The complex singularity exponent at o is defined to be*

$$c_o(\varphi) := \sup\{c \geq 0 : \mathcal{I}(c\varphi)_o = \mathcal{O}_o\}.$$

Our main result is the following:

Theorem 1.3. *Let φ be a plurisubharmonic function on $D \subset\subset \mathbb{C}^n$ with coordinates $z = (z_1, \cdots, z_n)$ containing the origin. Then for any $k \in \mathbb{N}$, there exist plurisubharmonic functions φ_k near the origin of $\mathbb{C}^n \times \mathbb{C}^{(2^k-1)n}$ with coordinates (z, w) such that*

Date: March 21, 2016.

Key words and phrases. L^2 extension theorem, plurisubharmonic function, Lelong number, multiplier ideal sheaf.

The authors were partially supported by NSFC-11431013. The second author would like to thank NTNU for offering him Onsager Professorship. The first author was partially supported by NSFC-11522101.

(1) $\varphi_k(z,o)=\varphi(z)$;
(2) $\nu(\varphi_k,(z,o))=\nu(\varphi,z)$;
(3) $\frac{(2^k-1)n}{\nu(\varphi,z)}\leq c_{(z,o)}(\varphi_k)\leq\frac{2^kn}{\nu(\varphi,z)}$
for any $z\in D$, where o is the origin in $\mathbb{C}^{(2^k-1)n}$.

The second inequality in (3) can be directly deduced by Skoda's result in [21] combined with (2).

Remark 1.4. *It is clear that* (3) *in Theorem 1.3 is equivalent to*

$$\frac{(2^k-1)n}{2^kn}\nu^{-1}(\varphi,z)\leq c_{(z,o)}((2^kn)\varphi_k)\leq\nu^{-1}(\varphi,z),$$

which implies

$$\{\nu(\varphi,z)\geq\frac{(2^k-1)n}{2^kn}c\}\supseteq\{c_{(z,o)}((2^kn)\varphi_k)\leq\frac{1}{c}\}\supseteq\{\nu(\varphi,z)\geq c\}.$$

Note that $\lim_{k\to+\infty}\frac{(2^k-1)n}{2^kn}=1$, *then we obtain*

$$\{z|\nu(\varphi,z)\geq c\}=\cap_k(\{z|c_{(z,o)}((2^kn)\varphi_k)\leq\frac{1}{c}\}).$$

By Berndtsson's solution of openness conjecture [1] posed by Demailly and Kollar [6] (for a proof in two dimensional case see [9, 8, 7]), it follows that $(\{z|c_{(z,o)}((2^kn)\varphi_k)\leq\frac{1}{c}\})$ is analytic for any $k\in\mathbb{N}$ and $c>0$, which deduces the following Siu's semi-continuity theorem for Lelong numbers [19] (For Demailly's new proof, the reader is refereed to [5] and [4])

Corollary 1.5. [19] $\{z|\nu(\varphi,z)\geq c\}$ *is an analytic set.*

We would like to thank the referee for pointing out that Berndtsson's solution of openness conjecture used above to deduce Corollary 1.5 might be replaced by the Hömander-Bombieri theorem as in Kiselman's proof of Siu's theorem.

2. Preparation

2.1. **Restriction formula about complex singularity exponent and Lelong number.** Let φ be a plurisubharmonic function on a neighborhood of the origin $o\in\mathbb{C}^n$. In [6], the following restriction formula ("important monotonicity result") about complex singularity exponents is obtained by using Ohsawa-Takegoshi L^2 extension theorem.

Proposition 2.1. [6] *For any regular complex submanifold $(H,o)\subset(\mathbb{C}^n,o)$,*

$$c_{o'}(\varphi|_H)\leq c_o(\varphi)\tag{2.1}$$

holds, where $\varphi|_H\not\equiv-\infty$, and o' emphasizes that $c_{o'}(u|_H)$ is computed on the submanifold H.

We recall the following restriction property of Lelong numbers.

Lemma 2.2. *(see* [5]*) For any regular complex submanifold $(H,o)\subset(\mathbb{C}^n,o)$,*

$$\nu(\varphi|_H,o')\geq\nu(\varphi,o)\tag{2.2}$$

holds, where $\varphi|_H\not\equiv-\infty$, and o' emphasizes that $\nu(\varphi|_H,o')$ is computed on the submanifold H.

2.2. **Lelong number and complex singularity exponent for $U(n)$ invariant plurisubharmonic functions on $\mathbb{C}^n$.** We recall the following characterization of $U(n)$ invariant plurisubharmonic function (see Lemma III.7.10) in [5]).

Lemma 2.3. *(see [5]) Let φ be a plurisubharmonic function on $\mathbb{B}^n$ which is $U(n)$ invariant. Then $\varphi = \chi(\log|z|)$, where χ is a convex increasing function.*

The folllowing remark is a direct consequence of Lemma 2.3

Remark 2.4. *(see [5]) Let φ be a plurisubharmonic function on $\mathbb{B}^n$ which is $U(n)$ invariant. Then $c_o(\varphi) = \frac{n}{\nu(\varphi,o)}$.*

Proof. By Definition 1.1, it is clear that

$$\nu(\varphi, o) = \lim_{t\to-\infty} \frac{\chi(t)}{t},$$

i.e. for any $\varepsilon > 0$, there exists $\delta > 0$ such that for any $|z| < \delta$,

$$(\nu(\varphi, o) + \varepsilon)\log|z| \leq \varphi(z) \leq (\nu(\varphi, o) - \varepsilon)\log|z|,$$

which deduces the present remark by directly calculating. □

2.3. **A holomorphic map and Lelong number.** Define a holomorphic map p_m from $\mathbb{C}^m \times \mathbb{C}^m$ to $\mathbb{C}^m$ with coordinates $(z_1, \cdots, z_m, w_1, \cdots, w_m)$ and $(\tilde{z}_1, \cdots, \tilde{z}_n)$ such that

$$p_m(z_1, \cdots, z_m, w_1, \cdots, w_m) = (z_1 - w_1, \cdots, z_m - w_m).$$

Let φ be a plurisubharmonic function on $D \subset \mathbb{C}^m$, then $p_m^*(\varphi)$ is well-defined on $p_m^{-1}(D) \subset \mathbb{C}^m \times \mathbb{C}^m$.

Lemma 2.5. *The construction of $p_m^*(\varphi)$ implies that for any $z_0 \in D$ the following statements hold:*

(1) $p_m^*(\varphi)(z_0, o) = \varphi(z_0)$;

(2) $\nu(p_m^*(\varphi), (z_0, o)) = \nu(\varphi, z_0) = \nu(p_m^*(\varphi)|_{z=z_0}, (z_0, o)')$, *where $(z_0, o)'$ emphasizes that $\nu(p_m^*(\varphi)|_{z=z_0}, (z_0, o)')$ is computed on the submanifold $\{z = z_0\}$.*

Proof. By definition of p_m, it follows that (1) and the second equality of (2) hold.

By (1), it follows that $\nu(p_m^*(\varphi), (z_0, o)) \leq \nu(\varphi, z_0)$ by Lemma 2.2.

It suffices to consider the case $z_0 = (0, \cdots, 0) \in D$. It is clear that $p_m^* \log|\tilde{z}| = \log|z - w| \leq \log(|z| + |w|)$. Then, for any $c > 0$ satisfying $\varphi \leq c\log|\tilde{z}| + O(1)$ $(z \to 0)$, we have $p_m^*\varphi \leq cp_m^* \log|\tilde{z}| + O(1) \leq c\log(|z| + |w|) + O(1)$, which implies $\nu(p_m^*(\varphi), (z_0, o)) \geq \nu(\varphi, z_0)$.

The lemma is proved. □

2.4. **An invariant property of Lelong numbers.** Let φ be a plurisuharmonic function on $\Omega \subset \mathbb{C}^n \times \mathbb{C}^m$ containing the origin, and let (z, w) denote the coordinates. One can define

$$\tilde{\varphi}(z, w) := \sup_{g\in U(m)} \varphi(z, gw)$$

on a $U(m)$ invariant neighborhood of $\Omega \cap \{w = o\}$. It is a plurisubharmonic function.

By definition 1.1, it follows that

Lemma 2.6. *One has $\nu(\tilde{\varphi}, (z_0, o)) = \nu(\varphi, (z_0, o))$ and $\nu(\tilde{\varphi}|_{z=z_0}, (z_0, o)') = \nu(\varphi|_{z=z_0}, (z_0, o)')$ hold for any $(z_0, o) \in (\Omega \cap \{w = o\})$.*

3. proof of Theorem 1.3

Let $m \sim 2^j n$ $(j = 0, 1, \cdots, k-1)$ as in Lemma 2.5, where $\sim$ means that the former is replaced by the latter. We consider the plurisubharmonic function $p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi)$ on domain in $\mathbb{C}^{2^k n}$.

By Lemma 2.5, it follows that

(1) $p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi)(z, o) = \varphi(z)$;

(2) $\nu(p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi), (z_0, o)) = \nu(\varphi, z_0) = \nu(p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi)|_{\{z=z_0\}}, (z_0, o)')$

for any $z_0 \in D$ and origin $o \in \mathbb{C}^{(2^k-1)n}$.

Let $\varphi_k(z, w) := \sup_{g \in U((2^k-1)n)}(p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi)(z, gw))$.

By Lemma 2.6 (the first and third equalities) and (2) (the third and fourth equalities), it follows that

$$\begin{aligned} \nu(\varphi_k|_{\{z=z_0\}}, (z_0, o)') &= \nu(p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi)|_{\{z=z_0\}}, (z_0, o)') \\ &= \nu(p^*_{2^{k-1}n} \circ \cdots \circ p^*_n(\varphi), (z_0, o)) \\ &= \nu(\varphi_k, (z_0, o)) = \nu(\varphi, z_0). \end{aligned} \tag{3.1}$$

By Remark 2.4 ($\mathbb{C}^n \sim \{z = z_0\} \cap \mathbb{C}^{2^k n} (= \{(z_0, w) : w \in \mathbb{C}^{(2^k-1)n}\})$, $\varphi \sim \varphi_k|_{\{z=z_0\}}$, $n \sim (2^k - 1)n$), it follows that

$$\nu(\varphi_k|_{\{z=z_0\}}, (z_0, o)') = \frac{(2^k - 1)n}{c_{(z_0,o)'}(\varphi_k|_{\{z=z_0\}})}. \tag{3.2}$$

From Proposition 2.1, equality 3.2 and equality 3.1, it follows that

$$c_{(z_0,o)}(\varphi_k) \geq c_{(z_0,o)'}(\varphi_k|_{\{z=z_0\}}) = \frac{(2^k - 1)n}{\nu(\varphi_k|_{\{z=z_0\}}, (z_0, o)')} = \frac{(2^k - 1)n}{\nu(\varphi, z_0)}. \tag{3.3}$$

Thus we finish proving the present theorem.

References

[1] B. Berndtsson, The openness conjecture for plurisubharmonic functions, arXiv:1305.5781.

[2] J.-P, Demailly, Nombres de Lelong généralisés, théorèmes d'intégralité et d'analyticité. (French) [Generalized Lelong numbers, integrality and analyticity theorems] Acta Math. 159 (1987), no. 3–4, 153–169.

[3] J.-P. Demailly, Multiplier ideal sheaves and analytic methods in algebraic geometry. School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000), 1–148, ICTP Lect. Notes, 6, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001.

[4] J.-P. Demailly, Analytic Methods in Algebraic Geometry, Higher Education Press, Beijing, 2010.

[5] J.-P. Demailly, Complex analytic and differential geometry, electronically accessible at http://www-fourier.ujf-grenoble.fr/ demailly/books.html.

[6] J-P. Demailly, J. Kollár, Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds. Ann. Sci. École Norm. Sup. (4) 34 (2001), no. 4, 525–556.

[7] C. Favre and J. Jonsson, The valuative tree. Lecture Notes in Mathematics, 1853. Springer-Verlag, Berlin, 2004. xiv+234 pp. ISBN: 3-540-22984-1.

[8] C. Favre and M. Jonsson, Valuative analysis of planar plurisubharmonic functions, Invent. Math. 162 (2005), no. 2, 271–311.

[9] C. Favre and M. Jonsson, Valuations and multiplier ideals, J. Amer. Math. Soc. 18 (2005), no. 3, 655–684.

[10] Q.A. Guan and X.Y. Zhou, Optimal constant in an L^2 extension problem and a proof of a conjecture of Ohsawa, SCIENCE CHINA Mathematics, January 2015 Vol. 58 No. 1: 35–59, doi: 10.1007/s11425-014-4946-4.

[11] Q.A. Guan and X.Y. Zhou, A solution of an L^2 extension problem with optimal estimate and applications, Ann. of Math. Volume 181 (2015), no. 3, 1139–1208.

[12] Q.A. Guan and X.Y. Zhou, A proof of Demailly's strong openness conjecture, Ann. of Math. (2) 182 (2015), no. 2, 605–616.
[13] Q.A. Guan and X.Y. Zhou, Effectiveness of Demailly's strong openness conjecture and related problems, Invent. Math. 202 (2015), no. 2, 635–676.
[14] Q.A. Guan and X.Y. Zhou, Characterization of multiplier ideal sheaves with weights of Lelong number one, Adv. Math. 285 (2015), 1688–1705.
[15] C.O. Kiselman, Un nombre de Lelong raffiné. Séminaire d'Analyse Complexe et Géométrie 1985–87, pp. 61–70. Faculté des Sciences de Tunis & Faculté des Sciences et Techniques de Monastir.
[16] C.O. Kiselman, Plurisubharmonic functions and potential theory in several complex variables. Development of mathematics 1950–2000, 655–714, Birkhäauser, Basel, 2000.
[17] J. Kollár (with 14 coauthors): Flips and Abundance for Algebraic Threefolds; Astérisque Vol. 211 (1992).
[18] V. Shokurov: 3-fold log flips; Izv. Russ. Acad. Nauk Ser. Mat. Vol. 56 (1992) 105–203.
[19] Y.T. Siu, Analyticity of sets associated to Lelong numbers and the extension of closed positive currents. Invent. Math. 27 (1974), 53–156.
[20] Y.T. Siu, Multiplier ideal sheaves in complex and algebraic geometry. Sci. China Ser. A 48 (2005), suppl., 1–31.
[21] H. Skoda, Sous-ensembles analytiques d'ordre fini ou infini dans $\mathbb{C}^n$, Bull. Soc. Math. France 100 (1972) 353–408.
[22] G. Tian, On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$, Invent. Math. 89 (1987), no. 2, 225–246.

Qi'an Guan: School of Mathematical Sciences, and Beijing International Center for Mathematical Research, Peking University, Beijing 100871, China.
E-mail address: guanqian@amss.ac.cn

Xiangyu Zhou: Institute of Mathematics, AMSS, and Hua Loo-Keng Key Laboratory of Mathematics, Chinese Academy of Sciences, Beijing 100190, China
E-mail address: xyzhou@math.ac.cn